# Earth System Model Evaluation with Observations to Constrain Future Climate Projections

Knowledge for Tomorrow

#### Veronika Eyring

Deutsches Zentrum für Luft- und Raumfahrt e.V. (DLR), Institute of Atmospheric Physics, Oberpfaffenhofen, Germany

23 November 2015 Math colloquium College of Engineering, Mathematics and Physical Sciences University of Exeter, UK



# Outline

#### 1. Introduction

- 2. How do we gain confidence in climate model projections?
- 3. Brief overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) design

#### 4. Opportunities for CMIP6

- More routine evaluation of Earth system models with observations
- Better consideration of internal variability and more process-oriented evaluation
- Emergent constraints: Use of observations to constrain a simulated future Earth system feedback

#### 5. Summary





DLR.de · Chart 3

# **1. Introduction**

# **The Climate System**



# How to improve long-term climate projections?





# **Historical Evolution of Climate Models**

- From Atmosphere-Ocean General Circulation Models (AOGCM) to Earth System Models (ESMs) -



# Coupled Model Intercomparison Project (CMIP)

- Understanding past, present and future climate -

- CMIP is a project of the World Climate Research Programme (WCRP)'s Working Group of Coupled Modelling (WGCM).
- Since 1995, CMIP has coordinated climate model experiments involving multiple international modeling teams worldwide.
- CMIP has led to a better understanding of past, present and future climate change and variability in a multi-model framework.
- CMIP defines common experiment protocols, forcings and output.
- CMIP has developed in phases, with the simulations of the fifth phase, CMIP5, now completed, and the planning of the sixth phase, i.e. CMIP6, well underway.
- CMIP's central goal is to advance scientific understanding of the Earth system.
- CMIP model simulations have also been regularly assessed as part of the IPCC Climate Assessments Reports and various national assessments.

# Why using an Ensemble of Model Simulation?

The multi-model approach is now a standard technique to assess projections of specific variables and to derive robust process understanding of the Earth's climate system in combination with observations.

- 1. Assessing the mechanisms responsible for model differences in poorly understood feedbacks
- 2. Estimating projection uncertainty
- 3. Determining why similarly forced models produce a range of responses.



IPCC, Figure SPM.7a, 2014





Possible temperature responses in 2081-2100 to low emission scenario RCP2.6



IPCC FAQ 12.1, Figure 1

# Human influence on the climate system is clear

Evidence of human influence has grown since the AR4.

It is *extremely likely* that human influence has been the **dominant cause of the observed warming since the mid-20th century**.

Figure SPM.6

IPCC AR5 Working Group I Climate Change 2013: The Physical Science Basis **Slide 9** 



# 2. How do we gain confidence in climate model projections?

## How do we gain confidence in climate model projections?

- Based on physical understanding of the climate system and its representation in • climate models, and
- On a demonstration of how well models represent a wide range of processes and ٠ climate characteristics on various spatial and temporal scales



#### 0.5

**Relative error measures** ot CMIP5 model performance (normalized by the median error of all model results), based on the global seasonalcycle climatology (1980-2005)

Climate models have continued to be developed and improved since the AR4.

**IPCC AR5 Working Group I** 

Slide 11

Chapter 9, Fig. 9.7

Climate Change 2013: The Physical Science Basis Following Gleckler et al. (2008) INTERGOVERNMENTAL PANEL ON Climate C



```
DLR.de · Chart 12
```

# A lot of progress has been made, but...

#### **Observations:**

- In many cases the lack or insufficient quality of long-term observations or observations for process evaluation remains an impediment.
- For many observational datasets formal error estimates are lacking.
- Disagreement in observations complicate model evaluation (e.g. in cloud property trends), and upper tropospheric / lower stratosphere (UTLS) temperature trends

**Systematic Biases:** e.g., Double Intertropical Convergence Zone (ITCZ), i.e. spurious ITCZ in the SH associated with excessive tropical precipitation or the equatorward bias in the SH atmospheric jet location



# **Equilibrium Climate Sensitivity Remains Uncertain**

#### Equilibrium climate sensitivity (ECS):

- Response of the climate system to constant RF on multi-century time scales.
- Defined as the change in global mean surface temperature at equilibrium that is caused by a doubling of the atmospheric CO2 concentration.



The model spread in ECS ranges from **2.1°C to 4.7°C** and is very similar to the assessment in AR4.

#### => Due to uncertainties in climate feedbacks



DLR.de · Chart 14

# 3. CMIP6 Design

# **CMIP Continuity**



A common suite of experiments for each phase of CMIP provides an opportunity to construct a multi-model ensemble using model output from various phases of CMIP



Eyring et al., GMD, in prep., 2015

# **CMIP: a More Continuous and Distributed Organization**

#### (3) CMIP-Endorsed Model Intercomparison Projects (MIPs)



#### (1) A handful of common experiments

#### DECK (entry card for CMIP)

- i. AMIP simulation (~1979-2014)
- ii. Pre-industrial control simulation
- iii. 1%/yr CO<sub>2</sub> increase
- iv. Abrupt 4xCO<sub>2</sub> run

#### CMIP6 Historical Simulation (entry card for CMIP6)

v. Historical simulation using CMIP6 forcings (1850-2014)

# (2) Standardization, coordination, infrastructure, documentation

DECK (Diagnosis, Evaluation, and Characterization of Klima Experiments) & CMIP6 Historical Simulation to be run for each model configuration used in CMIP6-Endorsed MIPs

Eyring et al., GMD, in prep., 2015

# Models are increasing in complexity and resolution

70s

From AOGCMs to Earth System Models with biogeochemical cycles, from lowres to highres

#### **Ozone layer** 130 km resolution orography **Upper-level** winds 1810 1610 1410 1210 1010 810 ioil moisture and temperatu 610 410 210 Surface winds 10 Sea ice Ocean bottom 25 km resolution orography topograph Growth of Climate Modeling **Coupled Climate Model** Atmospheric/Land Surface/Vegetation Sulfate Aerosol Ocean Sea Ice **Biogeochemical Cycles**

60s

https://www2.ucar.edu/news/understanding-climate-change-multimedia-gallery

9()c

80s

Solar energy

**Atmospheric Chemistry** 

Air-sea

Ocean currents,

Other components Atmospheric chemistry

Evaporation

Outgoing heat

**Upper Atmosphere** Atmospheric Chemistry

Marine Ecosystems

10s

temperature and salinity

exchanges

Cloud

Marine ecosyste

Dust/Sea Spray/Carbon Aerosols

**Ice Sheet** 

Interactive Vegetation

**Carbon Cycle** 

Vertical

overturni

# **4. Opportunities for CMIP6**

(A) More routine evaluation of Earth system models with observations

### **Earth System Model Evaluation**





#### **Routine Benchmarking and Evaluation Central Part of CMIP6**

CMIP evaluation tool to produce well-established analyses as soon as model output becomes available e.g., Community-develoed ESMValTool (Eyring et al., GMDD, 2015) and PCMDI metrics package (Gleckler et al., EOS, in press) - *Link to WGNE/WGCM Climate Model Metrics and Diagnostic Panel* 



Under-Exploited Observations for Model Evaluation Observations for Model Intercomparison Projects (obs4MIPs) WDAC Task Team on Observations for Model Evaluation

#### CMIP6





How to bring as much observational scrutiny as possible to the CMIP/IPCC process?

How to best utilize the wealth of satellite observations for the CMIP/IPCC process?

- Obs4MIPs has defined a set of technical specifications and criteria for developing observational data sets that are technically aligned with CMIP model output (with common file format, data and metadata structure).
- Over 50 datasets that conform to these standards are now archived on the ESGF alongside CMIP model output (<u>Teixeira et al., 2014</u>), including ESA CCI data
- Data users have enthusiastically received Obs4MIPs

#### **Routine Benchmarking and Evaluation Central Part of CMIP6**

- The objective is to enable routine model evaluation and to aid the individual modelling groups in their model development process by providing feedback concerning their model errors, particularly the systematic model errors.
- Building such a community-based capability is not meant to replace how CMIP research is currently performed but rather to complement it.



Higher-Order Characteristics of Models: State & Process Evaluation as Part of the Publication Process

Documentation of Progress over Time

•>

# 4. Opportunities for CMIP6

(B) Better consideration of internal variability and more process-oriented evaluation

### **Evaluation of Climate Models: Temperature trends**



Models reproduce observed temperature trends over many decades, including the more rapid warming since the mid-20th century and the cooling immediately following large volcanic eruptions (*very high confidence*).

Chapter 9, Fig. 9.8

1998–2012: 0.04 °C/decade 1951–2012: 0.11 °C/decade

#### Temperature Plateau ("Hiatus")

Internal decadal variability causes to a substantial degree **the difference between observations and the simulations**. There may also be a contribution from forcing inadequacies and, in some models, an overestimate of the response to increasing GHGs and other anthropogenic forcing (dominated by aerosols).



# Observed and simulated GMST trends in °C per decade

Box TS.3, Figure 1 Box 9.2, Figure 1 Noting Karl et al., 2015 and Trenberth. 2015





INTERGOVERNMENTAL PANEL ON Climate chan

### **Evaluation of Climate Models: Sea-Ice**



Robust evidence that the downward trend in Arctic summer sea-ice extent is better simulated than at the time of the AR4, with about one-quarter of the simulations showing a trend as strong as, or stronger, than in observations over the satellite era (since 1979).

IPCC AR5, Fig. 9.24 redone with ESMValTool

More than one ensemble member required to make robust assessments of model performance for a single model,

#### CMIP6-Endorsed Model Intercomparison Project HighResMIP Co-chairs: Rein Haarsma & Malcolm Roberts



# **4.** Opportunities for CMIP6

(C) Emergent constraints: Use of observations to constrain a simulated future Earth system feedback

# **Uncertainties in Projections of Future Climate**

#### 1. Internal Variability

- Due to the chaotic nature of climate system
- Noise of climate record is constant with time

#### 2. Emission Uncertainty

- Dominant uncertainty for long term projections estimated as mean of different scenarios
- Varying greenhouse gas emissions
- Land use change

#### 3. Climate Response Uncertainty

- Models are build on same principles but parametrizations are needed
- Increases when process become more relevant
- Decreases with model improvements and observational constraints





# **Emergent Constraints (ECs)**

- ECs are a relationship across an ensemble of models, between some aspect of Earth system sensitivity and an observable trend or variation in the current climate
  - > Emergent because it emerges from the ensemble of ESMs.
  - Constraint because it enables an observation to constrain the estimate of the Earth System sensitivity in the real world.
- The goal is to find a observable physical explanation to constrain the unobservable Earth system sensitivity



### **Emergent Constraint: Seasonal Cycle Physical Climate**



 $SAF \propto \Delta \alpha_s / \Delta T_s$ 

Large intermodel variations in the strength of snow albedo feedback (SAF) in climate change in the NH in April are nearly perfectly correlated with comparably large intermodel variations in feedback strength in the context of the seasonal cycle.

Hall and Qu, GRL, 2006

Feedback strength in the real seasonal cycle can be **observed** and compared to models.

These mostly fall outside the range of the observed estimate, suggesting many models have an unrealistic snow albedo feedback in the seasonal cycle context.



## How will the carbon cycle change with climate change?

Long-term change in land carbon uptake  $\Delta C_{Land}$  [GtC] (Friedlingstein et al., 2006):

 $\Delta C_{Land} = \gamma_{Long Term} \Delta T + \beta_{Long Term} \Delta C_{Atm}$ 

 $\Delta C_{Atm}$ : change in atm.  $CO_2$  concentration  $\Delta T$ : change in temperature  $\gamma, \beta$  : Feedback parameter

Response to increasing temperatures? Climate warming reduces the efficiency of CO<sub>2</sub> absorption by the land and ocean => more emitted carbon stays in the atmosphere leading to additional warming, representing a positive climate-carbon cycle feedback

The **carbon cycle-climate feedback**  $\gamma_{LT}$  quantifies the difference of (tropical) land carbon storage between coupled and uncoupled simulations due to climate change in terms of carbon loss per degree [GtC/K]; ("LT" = long term) :

$$\nu_{LT} = \frac{\Delta C_{LT}^c - \Delta C_{LT}^u}{\Delta T_T^c}$$

**Study 1** constrains  $\gamma_{LT}$ 

Cox et al., 2013; Wenzel, Cox, Eyring, Friedlingstein, JGR, 2014

 Response to increasing atmospheric CO2 concentrations? elevated CO2 will enhance Gross Primary Productivity (GPP)

The **carbon cycle-CO**<sub>2</sub> **concentration feedback**  $\beta_{LT}$  quantifies the change in land carbon due to the change in atmospheric CO<sub>2</sub> in the uncoupled simulation:  $\beta_{LT} = \frac{\Delta C_{LT}^u}{\Delta C^u}$ 

Study 2 constrains  $\beta_{LT}$ Wenzel, Cox, Eyring, Friedlingstein, in rev., 2015

#### Diagnosing the Carbon Cycle-Climate Feedback ( $\gamma_{LT}$ ) Coupled (1%COU) and uncoupled (1%BGC) 1%/yr simulations to estimate $\Delta C_{L}$



- Increasing temperature leads to decreased uptake of land carbon uptake
- Leads to higher CO<sub>2</sub> concentrations in the atmosphere
- Large uncertainties in the amount of future tropical land carbon uptake

Wenzel et al., JGR, 2014



# **Atmospheric CO<sub>2</sub> Measurements**

- Atmospheric CO<sub>2</sub> concentration increased over the last 50 years by approx. 100 ppmv
- Increasing atmospheric CO<sub>2</sub> concentration (black) mainly due to anthropogenic CO<sub>2</sub> emissions
- Seasonal variability (red) due to seasonal carbon cycle
  - Summer: more photosynthesis => atmospheric CO<sub>2</sub> decreases stored in terrestrial ecosystem
  - Winter: CO2 release by decomposition of soil organic matter





#### **Relationship Interannual Variations in Temperature & CO<sub>2</sub> Growth Rate**



- Responses of the carbon cycle to climate anomalies are mirrored in the interannual variability (IAV)
- Especially valid in the tropics, where strong variability caused by El Niño gives a spatially coherent pattern of warmer and colder years
  - Decomposition is mainly controlled by climate, warming for example increases microbial activity and therefore decomposition.
- > Observed relationship:  $\gamma_{IAV} = -4.4 \pm 0.9$  GtC/yr/K
- Same analysis for the CMIP5 ESMs using historical simulations

```
Wenzel et al., JGR, 2014
```

# Constraining the Carbon Cycle-Climate Feedback ( $\gamma_{LT}$ )



- EC between the long-term sensitivity of tropical land carbon storage to climate warming ( $\gamma_{LT}$ ) and the short-term sensitivity of atmospheric carbon dioxide (CO2) to interannual temperature variability ( $\gamma_{IAV}$ )
- Constrained carbon cycle-climate feedback parameter: -44 ± 14 GtC/K (unconstr. -49 ± 40 GtC/K)
- > Less carbon will be stored as a result of increasing of temperature
- > This leads to higher CO<sub>2</sub> concentrations in the atmosphere

Wenzel et al., JGR, 2014



# **Applying MDER to future Austral Jet Stream Positions**

- Multiple Diagnostic Ensemble Regression (MDER, Karpechko et al., 2013)
  - Uses process-oriented present day diagnostics to constrain future austral jet stream positions
- Equatorward bias of the CMIP5 models austral jet positions with a spread of 10°
- MDER is targeted to constrain near-term (2015-2034) projections of the austral jet position, and selects the historical jet position as the most important of 20 diagnostics.
- The method essentially recognizes the equatorward bias in the past jet position, and provides a bias correction of about 1.5° southward to future projections.



Wenzel et al., J. Clim., in press



a) 2015-2034

DLR.de · Chart 36

#### Weighting Model Projections: Arctic Sea-Ice Process-based constraints can be used to reduce the spread of model projections



- First year during which the September Arctic sea ice extent falls below a certain threshold is highly correlated with the September sea ice extent and annual mean sea ice volume averaged over the past.
- First year during which the September Arctic sea ice extent falls below a certain threshold are correlated with the past trend in September Arctic sea ice extent and the amplitude of the mean seasonal cycle of sea ice extent.
- Suggests a faster rate of summer Arctic sea ice decline than the multi-model mean
- A model is retained if, for each diagnostic, either this interval overlaps a ±20% interval around the observed/reanalysed value of the diagnostic or at least one ensemble member from that model gives a value for the diagnostic that falls within ±20% of the observational/reanalysed data.



### **Other Examples on Selected Feedbacks and ECS**

- The extratropical surface cryosphere feedback was constrained by Crook & Forster (2014) using variations in the seasonal cycle of the cryosphere. Models were found to largely underestimate this feedback (0.4 – 1.2 W m<sup>-2</sup> K-1 compared to 3.1 ± 1.3 W m<sup>-2</sup> K<sup>-1</sup>) under warming despite their comparable seasonal sensitivity to observations.
- Gordon et al. (2013) related the water vapor feedback to observed variability (2002-2009)
  - Demonstrated the physical explanation of the relation between short and long-term forced changes in models under warming.
  - However, relative weak relation combined with large uncertainties in the observations.
  - Suggested an observational record of 25 years or longer could significantly improve the demonstrated observational constraint.
- Tian (2015) shows that show that the double-ITCZ bias and ECS in 44 GCMs from CMIP 3/5 are negatively correlated
  - Southern ITCZ index: model climatological annual mean precip bias over southeastern Pacific
  - Low sensitivity models having problems in representing its southern branch.
  - ECS might be in the higher end of its range (~4.0°C) and most CMIP3/5 models might have underestimated ECS.



# **Constraining Equilibrium Climate Sensitivity**



- Spread in ECS arises largely from low clouds
- Relates ECS to the strength of mixing in the lower troposphere over warm tropical oceans (lower-tropospheric mixing index LTMI)
- Higher-sensititivity models simulate certain cloud-relevant phenomena better.
- However, the metric suffers from large uncertainties in the observed estimates.

 No final conclusions can be made yet, but the results of EC studies generally suggest an underestimation of ECS by models due to cryospheric and cloud feedbacks.



#### Implied Sensitivity



# Summary

While progress has been made in ESM evaluation over the last decades, there are important opportunities and challenges for CMIP6, with simulations starting in 2016

- In many cases the lack or insufficient quality of long-term observations or observations for process evaluation remains an impediment, but improvements can be made by fully exploiting existing observations and by taking into account observational uncertainty.
- Make the evaluation of CMIP models with well-established diagnostics and performance metrics more routine (by developing and applying diagnostic tools such as the ESMValTool) to leave more time for innovative research.
- Part of the difference between model results and observations can be attributed to unforced variability, originating from the nonlinear nature of the variable climate system. An accurate assessment of model performance therefore has to take into account internal climate variability in addition to observational uncertainty.



# **Summary Emergent Constraints**

- While evaluation of the evolving climate state and processes can be used to build confidence in model fidelity, this does not guarantee the correct response to changed forcing in the future.
- Emergent constraint analysis refers to the use of observations to constrain a simulated future Earth system feedback offers the potential to reduce uncertainty in climate projections.
- Studies have been published that focus both on constraining ECS more generally, but also on constraining individual key feedbacks at a process level.
- ECs studies can help guiding model development onto processes crucial to the magnitude and spread of future Earth system change. This can also be used to prioritize future observations activities.
- >A necessary property of emergent constraints is a physical basis for the relation.
- There are many open questions and issues, but emergent constraints remain a promising approach that should be fully exploited in CMIP analysis.